Neuroscience Letters, 164 (1993) 236–238
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Acuña-Castroviejo, D., see Gomar, M.D., 149
Akaike, A., see Shimohama, S., 55
Akiguchi, I., see Nagao, M., 221
Akiyama, H., Ikeda, K., Kondo, H., Kato, M. and McGeer, P.L., Microglia express the type 2 plasminogen activator inhibitor in the brain of control subjects and patients with Alzheimer's disease, 233
Alberici, A., see Valerio. A., 109
Albuquerque, E.X., see Castro, N.G., 137
Amano, M., see Kubo, T., 113
Amenta, F., see Bronzetti, E., 47
Anadón, R., see Yáñez, J., 213
Asari, T., see Kubo, T., 113

Banerjee, A., Martin, T.F.J. and DasGupta, B.R., Nerve growth factor induces sensitivity to botulinum neurotoxin type A in norepinephrine-secreting PC12 cells, 93
Bannon, M.J., see Whitty, C.J., 141
Bär, P.R., see Joosten, E.A.J., 85
Barnes, C.D., see Wang, Z., 117
Barone, F.C., see Liu, T., 125
Behrens, U.D., Douglas, R.H. and Wagner, H.-J., Gonadotropin-releasing hormone, a neuropeptide of efferent projections to the teleost retina induces light-adaptive spinule formation on horizontal cell dendrites in dark-adapted preparations kept in vitro, 59
Belloni, M., see Valerio, A., 109
Bennis, M., see M'hamed, S.B., 195
Berridge, C.W., see Page, M.E., 81
Bickford, P.C., see Lin, A.M.-Y., 71
Bing, O., Möller, C., Engel, J.A., Söderpalm, B. and Heilig, M., Anxiolytic-like action of centrally administered galanin, 17
Bourre, J.-M., see Clement, M., 163
Bowker, R.M., Sonea, I.M., Vex, K.B. and Caron, J.P., Substance P innervation of equine synovial membranes: joint differences and neural and nonneural receptor localizations, 76
Bregman, B.S., see Joosten, E.A.J., 85
Bronzetti, E., Felici, L. and Amenta, F., Effect of ipsilateral lesioning of the nucleus basalis magnocellularis and of L-α-glyceryl phosphorylcholine treatment on choline acetyltransferase and acetylcholinesterase in the rat fronto-parietal cortex, 47
Bryant, H.J., see Kowtha, V.C., 129
Buchan, A.M., see MacManus, J.P., 89
Buonamici, M., see Valerio, A., 109
Byers, M.R., see Oswald, R.J., 190

Camp, D.M., see Crippens, D., 145
Caron, J.P., see Bowker, R.M., 76
Casillas, T., Delicado, E.G. and Miras-Portugal, M.T., Adenosine 5′-triphosphate modulation of nitrobenzylthioinosine binding sites in plasma membranes of bovine chromaffin cells, 51
Castillo, J.L., see Gomar, M.D., 149
Castro, N.G. and Albuquerque, E.X., Brief-lifetime, fast-inactivating ion channels account for the α-bungarotoxin-sensitive nicotinic response in hippocampal neurons, 137
Chin, W.W., see Koibuchi, N., 159
Choi, D.W., see Hewett, S.J., 229
Ciriello, J., see Solano-Flores, L.P., 217
Clement, M. and Bourre, J.-M., Alteration of brain and liver microsomal polyunsaturated fatty acids following dietary vitamin E deficiency, 163
Clinton, J., Mann, D.M.A. and Roberts, G.W., Frontal lobe dementia is not a variant of prion disease, 1
Clow, A., Freestone, C., Lewis, E., Dexter, D., Sandler, M. and Glover, V., The effect of pergolide and MDL 72974 on rat brain CuZn superoxide dismutase, 41
Corbett, J.A., see Hewett, S.J., 229
Crippens, D., Camp, D.M. and Robinson, T.E., Basal extracellular dopamine in the nucleus accumbens during amphetamine withdrawal: a 'no net flux' microdialysis study, 145

DasGupta, B.R., see Banerjee, A., 93
Dash, P.K. and Moore, A.N., Inhibitors of endocytosis, endosome fusion, and lysosomal processing inhibit the intracellular proteolysis of the amyloid precursor protein, 183
De Backer, J.-P., see De Keyser, J., 63
De Keyser, J., Vauquelin, G., De Backer, J.-P., De Vos, H. and Wilczak, N., What intracranial tissues in humans contain sumatriptan-sensitive serotonin 5-HT_1-type receptors?, 63
De Vos, H., see De Keyser, J., 63
De Zeeuw, C.I., see Wentzel, P.R., 25
del Aguila, C.M., see Gomar, M.D., 149
Delicado, E.G., see Casillas, T., 51
Dexter, D., see Clow, A., 41
Díaz-Ruiz, C., Pérez-Tomás, R., Domingo, J. and Ferrer, I., Immunohistochemical localization of transforming growth factor-α in choroid plexus of the rat and chicken, 44
Ding, Y.-q., Wang, Y.-q., Qin, B.-z. and Li, J.-s., The major pelvic ganglion is the main source of nitric oxide synthase-containing nerve fibers in penile erectile tissue of the rat, 187
Domingo, J., see Díaz-Ruiz, C., 44
Douglas, R.H., see Behrens, U.D., 59

Ekström, P., see Yáñez, J., 213
Engel, J.A., see Bing, O., 17
E.Wisniewski, K., see Kida, E., 121

Feldman, A.G., see St-Onge, N., 171
Felici, L., see Bronzetti, E., 47
Fernández, B., see Gomar, M.D., 149
Ferrer, I., see Díaz-Ruiz, C., 44
Feuerstein, G.Z., see Liu, T., 125
Flavin, M.P., Yang, Y. and Ho, G., Hypoxic forebrain cholinergic neuron injury: role of glucose, excitatory amino acid receptors and nitric oxide, 5
Foote, S.L., see Page, M.E., 81
Franken, P., see Tobler, I., 105
Freestone, C., see Clow, A., 41
Freund, R.K., Wang, Y. and Palmer, M.R., Differential effects of ethanol on the firing rates of Golgi-like neurons and Purkinje neurons in cerebellar slices in vitro, 9
Fu, W.-M. and Lin, J.-L., Developmental change in the modulation of acetylcholine receptor channel by protein kinase C activation in *Xenopus* embryonic muscle cells, 97
Fung, S.J., see Wang, Z., 117

Neuroscience Letters, 164 (1993) 236–238
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Acuña-Castroviejo, D., see Gomar, M.D., 149
Akaike, A., see Shimohama, S., 55
Akiguchi, I., see Nagao, M., 221
Akiyama, H., Ikeda, K., Kondo, H., Kato, M. and McGeer, P.L., Microglia express the type 2 plasminogen activator inhibitor in the brain of control subjects and patients with Alzheimer's disease, 233
Alberici, A., see Valerio. A., 109
Albuquerque, E.X., see Castro, N.G., 137
Amano, M., see Kubo, T., 113
Amenta, F., see Bronzetti, E., 47
Anadón, R., see Yáñez, J., 213
Asari, T., see Kubo, T., 113

Banerjee, A., Martin, T.F.J. and DasGupta, B.R., Nerve growth factor induces sensitivity to botulinum neurotoxin type A in norepinephrine-secreting PC12 cells, 93
Bannon, M.J., see Whitty, C.J., 141
Bär, P.R., see Joosten, E.A.J., 85
Barnes, C.D., see Wang, Z., 117
Barone, F.C., see Liu, T., 125
Behrens, U.D., Douglas, R.H. and Wagner, H.-J., Gonadotropin-releasing hormone, a neuropeptide of efferent projections to the teleost retina induces light-adaptive spinule formation on horizontal cell dendrites in dark-adapted preparations kept in vitro, 59
Belloni, M., see Valerio, A., 109
Bennis, M., see M'hamed, S.B., 195
Berridge, C.W., see Page, M.E., 81
Bickford, P.C., see Lin, A.M.-Y., 71
Bing, O., Möller, C., Engel, J.A., Söderpalm, B. and Heilig, M., Anxiolytic-like action of centrally administered galanin, 17
Bourre, J.-M., see Clement, M., 163
Bowker, R.M., Sonea, I.M., Vex, K.B. and Caron, J.P., Substance P innervation of equine synovial membranes: joint differences and neural and nonneural receptor localizations, 76
Bregman, B.S., see Joosten, E.A.J., 85
Bronzetti, E., Felici, L. and Amenta, F., Effect of ipsilateral lesioning of the nucleus basalis magnocellularis and of L-α-glyceryl phosphorylcholine treatment on choline acetyltransferase and acetylcholinesterase in the rat fronto-parietal cortex, 47
Bryant, H.J., see Kowtha, V.C., 129
Buchan, A.M., see MacManus, J.P., 89
Buonamici, M., see Valerio, A., 109
Byers, M.R., see Oswald, R.J., 190

Camp, D.M., see Crippens, D., 145
Caron, J.P., see Bowker, R.M., 76
Casillas, T., Delicado, E.G. and Miras-Portugal, M.T., Adenosine 5′-triphosphate modulation of nitrobenzylthioinosine binding sites in plasma membranes of bovine chromaffin cells, 51
Castillo, J.L., see Gomar, M.D., 149
Castro, N.G. and Albuquerque, E.X., Brief-lifetime, fast-inactivating ion channels account for the α-bungarotoxin-sensitive nicotinic response in hippocampal neurons, 137
Chin, W.W., see Koibuchi, N., 159
Choi, D.W., see Hewett, S.J., 229
Ciriello, J., see Solano-Flores, L.P., 217
Clement, M. and Bourre, J.-M., Alteration of brain and liver microsomal polyunsaturated fatty acids following dietary vitamin E deficiency, 163
Clinton, J., Mann, D.M.A. and Roberts, G.W., Frontal lobe dementia is not a variant of prion disease, 1
Clow, A., Freestone, C., Lewis, E., Dexter, D., Sandler, M. and Glover, V., The effect of pergolide and MDL 72974 on rat brain CuZn superoxide dismutase, 41
Corbett, J.A., see Hewett, S.J., 229
Crippens, D., Camp, D.M. and Robinson, T.E., Basal extracellular dopamine in the nucleus accumbens during amphetamine withdrawal: a 'no net flux' microdialysis study, 145

DasGupta, B.R., see Banerjee, A., 93
Dash, P.K. and Moore, A.N., Inhibitors of endocytosis, endosome fusion, and lysosomal processing inhibit the intracellular proteolysis of the amyloid precursor protein, 183
De Backer, J.-P., see De Keyser, J., 63
De Keyser, J., Vauquelin, G., De Backer, J.-P., De Vos, H. and Wilczak, N., What intracranial tissues in humans contain sumatriptan-sensitive serotonin 5-HT_1-type receptors?, 63
De Vos, H., see De Keyser, J., 63
De Zeeuw, C.I., see Wentzel, P.R., 25
del Aguila, C.M., see Gomar, M.D., 149
Delicado, E.G., see Casillas, T., 51
Dexter, D., see Clow, A., 41
Díaz-Ruiz, C., Pérez-Tomás, R., Domingo, J. and Ferrer, I., Immunohistochemical localization of transforming growth factor-α in choroid plexus of the rat and chicken, 44
Ding, Y.-q., Wang, Y.-q., Qin, B.-z. and Li, J.-s., The major pelvic ganglion is the main source of nitric oxide synthase-containing nerve fibers in penile erectile tissue of the rat, 187
Domingo, J., see Díaz-Ruiz, C., 44
Douglas, R.H., see Behrens, U.D., 59

Ekström, P., see Yáñez, J., 213
Engel, J.A., see Bing, O., 17
E.Wisniewski, K., see Kida, E., 121

Feldman, A.G., see St-Onge, N., 171
Felici, L., see Bronzetti, E., 47
Fernández, B., see Gomar, M.D., 149
Ferrer, I., see Díaz-Ruiz, C., 44
Feuerstein, G.Z., see Liu, T., 125
Flavin, M.P., Yang, Y. and Ho, G., Hypoxic forebrain cholinergic neuron injury: role of glucose, excitatory amino acid receptors and nitric oxide, 5
Foote, S.L., see Page, M.E., 81
Franken, P., see Tobler, I., 105
Freestone, C., see Clow, A., 41
Freund, R.K., Wang, Y. and Palmer, M.R., Differential effects of ethanol on the firing rates of Golgi-like neurons and Purkinje neurons in cerebellar slices in vitro, 9
Fu, W.-M. and Lin, J.-L., Developmental change in the modulation of acetylcholine receptor channel by protein kinase C activation in *Xenopus* embryonic muscle cells, 97
Fung, S.J., see Wang, Z., 117

Gerhardt, G.A., see Lin, A.M.-Y., 71
Gerrits, N.M., see Wentzel, P.R., 25
Gibbs, R.B., see Koibuchi, N., 159
Gispen, W.H., see Joosten, E.A.J., 85
Glover, V., see Clow, A., 41
Göder, R., Häbler, H.-J., Jänig, W. and Michaelis, M., Receptive properties of afferent nerve fibres associated with the rat saphenous vein, 175
Golabek, A.A., see Kida, E., 121
Gomar, M.D., Fernández, B., Castillo, J.L., del Aguila, C.M. and Acuña-Castroviejo, D., Melatonin counteracts pinealectomy-dependent decreases in rat brain [^{3}H]flunitrazepam binding through an opioid mechanism, 149

Häbler, H.-J., see Göder, R., 175
Haro, R. and Paz, C., Effects of ozone exposure during pregnancy on ontogeny of sleep in rats, 67
Hashimoto, K., see Hayama, T., 13
Hayama, T., Hashimoto, K. and Ogawa, H., Projection of the inferior dental nerve to the primary somatosensory cortex in rats, 13
Heilig, M., see Bing, O., 17
Heinemann, U., see Strecker, T., 209
Hewett, S.J., Corbett, J.A., McDaniel, M.L. and Choi, D.W., Interferon-γ and interleukin-1β induce nitric oxide formation from primary mouse astrocytes, 229
Hill, I.E., see MacManus, J.P., 89
Ho, G., see Flavin, M.P., 5
Holmqvist, B.I., see Yáñez, J., 213
Holstege, J.C., see Wentzel, P.R., 25
Hopp, K., see Schulte-Bockholt, A., 101

Igwe, O.J. and Ning, L., Inositol 1,4,5-trisphosphate arm of the phosphatidylinositide signal transduction pathway in the rat cerebellum during aging, 167
Ikeda, K., see Akiyama, H., 233
Iseki, T., see Sato, J., 225
Iwakawa, Y., see Kohyama, J., 30

Jänig, W., see Göder, R., 175
Joëls, M., see Karst, H., 154
Jones, K.E., see Koibuchi, N., 159
Joosten, E.A.J., Bär, P.R., Gispen, W.H. and Bregman, B.S., Transient projections from rat occipital cortex are able to respond to a spinal target derived diffusible factor in vitro, 85

Kabasawa, H., see Ooka-Souda, S., 33
Kadota, T., see Ooka-Souda, S., 33
Kalcher, J., see Pfurtscheller, G., 179
Kamo, H., see Nagao, M., 221
Kapatos, G., see Whitty, C.J., 141
Karst, H., Joëls, M. and Wadman, W.J., Low-threshold calcium current in dendrites of the adult rat hippocampus, 154
Kato, M., see Akiyama, H., 233
Kida, E., E.Wisniewski, K. and Golabek, A.A., Increased expression of subunit c of mitochondrial ATP synthase in brain tissue from neuronal ceroid lipofuscinoses and mucopolysaccharidosis cases but not in long-term fibroblast cultures, 121
Kimura, J., see Nagao, M., 221
Kimura, J., see Shimohama, S., 55
Kleinhaus, A.L., see Sahley, C.L., 37
Koch, T.R., see Schulte-Bockholt, A., 101
Kohji, T., see Kohyama, J., 30
Kohyama, J., Kohji, T., Shimohira, M. and Iwakawa, Y., Phasic loss of intercostal muscle activity occurring with rapid eye movements during REM sleep, 30
Koibuchi, N., Gibbs, R.B., Jones, K.E., Yamaoka, S., Chin, W.W., Pfaff, D.W. and Suzuki, M., Increase in c-*erb*Aα2 mRNA in the parvocellular region of the paraventricular nucleus of the hypothalamus following thyroidectomy in the adult male rat, 159
Kondo, H., see Akiyama, H., 233
Kowtha, V.C., Quong, J.N., Bryant, H.J. and Stenger, D.A., Comparative electrophysiological properties of NG108–15 cells in serum-containing and serum-free media, 129
Kubo, T., Amano, M. and Asari, T., *N*-Methyl-D-aspartate receptors but not non-*N*-methyl-D-aspartate receptors mediate hypertension induced by carotid body chemoreceptor stimulation in the rostral ventrolateral medulla of the rat, 113
Kumazawa, T., see Sato, J., 225

Lewis, E., see Clow, A., 41
Li, J.-s., see Ding, Y.-q., 187
Lin, A.M.-Y., Bickford, P.C., Palmer, M.R. and Gerhardt, G.A., Ethanol inhibits the uptake of exogenous norepinephrine from the extracellular space of the rat cerebellum, 71
Lin, J.-L., see Fu, W.-M., 97
Liu, R.-H., see Wang, Z., 117
Liu, T., Young, P.R., McDonnell, P.C., White, R.F., Barone, F.C. and Feuerstein, G.Z., Cytokine-induced neutrophil chemoattractant mRNA expressed in cerebral ischemia, 125

M'hamed, S.B., Sequeira, H., Poulain, P., Bennis, M. and Roy, J.C., Sensorimotor cortex projections to the ventrolateral and the dorsomedial medulla oblongata in the rat, 195
MacManus, J.P., Buchan, A.M., Hill, I.E., Rasquinha, I. and Preston, E., Global ischemia can cause DNA fragmentation indicative of apoptosis in rat brain, 89
Maki, B.E., see McIlroy, W.E., 199
Mann, D.M.A., see Clinton, J., 1
Martin, T.F.J., see Banerjee, A., 93
McDaniel, M.L., see Hewett, S.J., 229
McDonnell, P.C., see Liu, T., 125
McGeer, P.L., see Akiyama, H., 233
McIlroy, W.E. and Maki, B.E., Do anticipatory postural adjustments precede compensatory stepping reactions evoked by perturbation?, 199
Meier, D.A., see Schulte-Bockholt, A., 101
Memo, M., see Valerio, A., 109
Meßlinger, K., Schepelmann, K., Pawlak, M. and Schmidt, R.F., Bradykinin B_1 and B_2 receptor antagonists do not change the ongoing activity of slowly conducting articular afferents in the inflamed knee joint of the cat, 21
Michaelis, M., see Göder, R., 175
Miras-Portugal, M.T., see Casillas, T., 51
Möller, C., see Bing, O., 17
Moore, A.N., see Dash, P.K., 183

Nagao, M., Oka, N., Kamo, H., Akiguchi, I. and Kimura, J., *Ulex europaeus* I and glycin max bind to the human olfactory bulb, 221
Neuper, C., see Pfurtscheller, G., 179
Ning, L., see Igwe, O.J., 167

Ogawa, H., see Hayama, T., 13
Ohara, O., see Shimohama, S., 55
Oka, N., see Nagao, M., 221

Ooka-Souda, S., Kadota, T. and Kabasawa, H., The preoptic nucleus: the probable location of the circadian pacemaker of the hagfish, *Eptatretus burgeri*, 33
Oswald, R.J. and Byers, M.R., The injury response of pulpal NPY-IR sympathetic fibers differs from that of sensory afferent fibers, 190
Otterson, M.F., see Schulte-Bockholt, A., 101

Page, M.E., Berridge, C.W., Foote, S.L. and Valentino, R.J., Corticotropin-releasing factor in the locus coeruleus mediates EEG activation associated with hypotensive stress, 81
Palmer, M.R., see Freund, R.K., 9
Palmer, M.R., see Lin, A.M.-Y., 71
Pawlak, M., see Meßlinger, K., 21
Paz, C., see Haro, R., 67
Pérez-Tomás, R., see Díaz-Ruiz, C., 44
Pfaff, D.W., see Koibuchi, N., 159
Pfurtscheller, G., Neuper, C. and Kalcher, J., 40-Hz oscillations during motor behavior in man, 179
Poulain, P., see M'hamed, S.B., 195
Preston, E., see MacManus, J.P., 89

Qi, H., see St-Onge, N., 171
Qin, B.-z., see Ding, Y.-q., 187
Quong, J.N., see Kowtha, V.C., 129

Rasquinha, I., see MacManus, J.P., 89
Reddy, V.K., see Wang, Z., 117
Roberts, G.W., see Clinton, J., 1
Robinson, T.E., see Crippens, D., 145
Rosas-Arellano, M.P., see Solano-Flores, L.P., 217
Roy, J.C., see M'hamed, S.B., 195

Sahley, C.L., Strong, J.A. and Kleinhaus, A.L., FMRF-amide modulates the electrical activity of the leech Retzius cell, 37
Sandler, M., see Clow, A., 41
Sato, J., Suzuki, S., Iseki, T. and Kumazawa, T., Adrenergic excitation of cutaneous nociceptors in chronically inflamed rats, 225
Schepelmann, K., see Meßlinger, K., 21
Schmidt, R.F., see Meßlinger, K., 21
Schulte-Bockholt, A., Meier, D.A., Hopp, K., Stadelmann, A., Otterson, M.F., Telford, G.L. and Koch, T.R., Gene expression and transcript size of the prepro-peptide VIP/PHM-27 in normal human tissue, 101
Sequeira, H., see M'hamed, S.B., 195
Shimohama, S., Tamura, Y., Akaike, A., Tsukahara, T., Ohara, O., Watanabe, S. and Kimura, J., Brain-derived neurotrophic factor pretreatment exerts a partially protective effect against glutamate-induced neurotoxicity in cultured rat cortical neurons, 55
Shimohira, M., see Kohyama, J., 30
Söderpalm, B., see Bing, O., 17
Solano-Flores, L.P., Rosas-Arellano, M.P. and Ciriello, J., C-fos expression in arcuate nucleus following intracerebroventricular hypertonic saline injections, 217
Sonea, I.M., see Bowker, R.M., 76
Spano, P.F., see Valerio, A., 109
Stadelmann, A., see Schulte-Bockholt, A., 101
Stenger, D.A., see Kowtha, V.C., 129
St-Onge, N., Qi, H. and Feldman, A.G., The patterns of control signals underlying elbow joint movements in humans, 171
Strecker, T. and Heinemann, U., Redistribution of K^+ channels into dendrites is unlikely to account for developmental down regulation of A-currents in rat dentate gyrus granule cells, 209
Strong, J.A., see Sahley, C.L., 37
Suzuki, M., see Koibuchi, N., 159
Suzuki, S., see Sato, J., 225

Taglietti, V., see Toselli, M., 134
Takahashi, J.S., see Zhang, Y., 203
Tamura, Y., see Shimohama, S., 55
Telford, G.L., see Schulte-Bockholt, A., 101
Tinti, C., see Valerio, A., 109
Tobler, I. and Franken, P., Sleep homeostasis in the guinea pig: similar response to sleep deprivation in the light and dark period, 105
Toselli, M. and Taglietti, V., Baclofen inhibits high-threshold calcium currents with two distinct modes in rat hippocampal neurons, 134
Tsukahara, T., see Shimohama, S., 55
Turek, F.W., see Zhang, Y., 203

Valentino, R.J., see Page, M.E., 81
Valerio, A., Tinti, C., Alberici, A., Belloni, M., Buonamici, M., Spano, P.F. and Memo, M., Deafferentation induces early and delayed differential changes in the pattern of expression of the various guanine nucleotide binding protein mRNAs in rat striatum, 109
Van Reeth, O., see Zhang, Y., 203
Vauquelin, G., see De Keyser, J., 63
Vex, K.B., see Bowker, R.M., 76

Wadman, W.J., see Karst, H., 154
Wagner, H.-J., see Behrens, U.D., 59
Wang, Y., see Freund, R.K., 9
Wang, Y.-q., see Ding, Y.-q., 187
Wang, Z., Liu, R.-H., Fung, S.J., Reddy, V.K. and Barnes, C.D., Immunohistochemical localization of glutamate-containing neurons in the lateral reticular nucleus projecting to the cerebellar vermis in the kitten, 117
Watanabe, S., see Shimohama, S., 55
Wentzel, P.R., De Zeeuw, C.I., Holstege, J.C. and Gerrits, N.M., Colocalization of GABA and glycine in the rabbit oculomotor nucleus, 25
White, R.F., see Liu, T., 125
Whitty, C.J., Kapatos, G. and Bannon, M.J., Neurotrophic effects of substance P on hippocampal neurons in vitro, 141
Wilczak, N., see De Keyser, J., 63

Yamaoka, S., see Koibuchi, N., 159
Yáñez, J., Anadón, R., Holmqvist, B.I. and Ekström, P., Neural projections of the pineal organ in the larval sea lamprey (*Petromyzon marinus* L.) revealed by indocarbocyanine dye tracing, 213
Yang, Y., see Flavin, M.P., 5
Young, P.R., see Liu, T., 125

Zee, P.C., see Zhang, Y., 203
Zhang, Y., Van Reeth, O., Zee, P.C., Takahashi, J.S. and Turek, F.W., Fos protein expression in the circadian clock is not associated with phase shifts induced by a nonphotic stimulus, triazolam, 203